

National Wastewater Surveillance Programme - COVID-19

Week 14 2022 (Week ending 10 April 2022)

Report prepared on 14 April 2022

Overview

SARS-CoV-2, the virus that causes COVID-19 disease, is shed in the faeces of people that are infected and so the viral RNA can be detected in wastewater. As such, testing wastewater for SARS-CoV-2 RNA is an efficient population-based COVID-19 surveillance tool. Based on national and international data, this method has been shown to be an indicator of increasing and decreasing cases (i.e., early warning system) and complements other surveillance tools. A national wastewater COVID-19 surveillance programme was established in 2021 by the Institute of Environmental Science and Research (ESR). This work is funded by the New Zealand Ministry of Health and is part of New Zealand's COVID-19 response.

Wastewater samples are collected from wastewater treatment plants across both the North and South islands of New Zealand. In addition, in Auckland there are a small number of sample sites that are pump stations and manholes. Most sites are sampled at least weekly between Monday and Thursday of any given week. The number of sites and frequency of collection varies over time. Grab or 24-hour composite samples are collected.

Approach

Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing involves the concentration of virus and extraction of viral RNA. The presence of SARS-CoV-2 RNA in the sample is then determined using RT-qPCR.

A result of '**not detected**', means that SARS-CoV-2 RNA is either absent from the sample, or at a level too low to be detected. When SARS-CoV-2 RNA is **detected**, the concentration in the sample can be calculated. Low amounts of SARS-CoV-2 RNA in a sample may not be able to be accurately quantified and are recorded as less than the limit of quantitation. For quantitation, the raw concentration data (i.e., genome copies per L or per mL) is converted to a viral load of **genome copies per day per person**. This calculation takes into account the flow rate of wastewater entering the wastewater treatment plant and the population in the catchment. This is the population-normalised viral load.

Key Points and Limitations

- SARS-CoV-2 RNA concentrations **should not be compared between wastewater catchments**.
- Day to day variability in SARS-CoV-2 RNA concentrations, especially in smaller catchments, is to be expected. Greater variability is expected with grab samples.
- Generally, increasing viral loads are associated with increasing numbers of people with SARS-CoV-2 infection and vice versa (decreasing concentrations indicating decreasing cases). However, there are a number of factors that affect the amount of viral RNA detected and so data from wastewater surveillance cannot indicate the exact number of COVID-19 cases in the catchment area.
- Results from wastewater testing should not be used in isolation but as an additional tool used for early detection and monitoring trends.
- The number of COVID-19 cases reported via individual testing are reported for each region to provide a comparison to the wastewater results. The cases in each catchment area are an estimate of the number of people in that wastewater catchment area that have reported a positive test. However, because the wastewater catchments do not exactly align with regional boundaries, the number of cases estimated by region and by water catchment area may be different.
- Data are provisional and may be subject to change by location.
- As septic tank systems are not connected to wastewater treatment plants, the wastewater from these households will not be represented in the data.

Please see notes at the end of the document for further details.

Results from Week 14 2022

In week 14, 2022 (week ending 10 April 2022), wastewater samples were received and analysed from 120 individual sites in New Zealand. SARS-CoV-2 RNA was detected in 119 of these sites. Figure 1 below gives the proportion of sites with positive results over time. Figure 2 is a map of locations sampled in the most recent week. Pages 6-26 provide overviews of SARS-CoV-2 levels from individual sites.

Table 1 provides weekly data from all sites from August 2021 (Week 32) to April 2022 (Week 14).




Figure 1. Wastewater SARS-CoV-2 (COVID-19) surveillance results from week 36, 2021 to week 14, 2022.





Figure 2. Location of sampling sites and latest results in the two weeks ending 10 April 2022.

Interpreting site graphs

Sample collection method. Results based on 24-hour composite samples, collected using an autosampler, may be more representative than from grab samples (see notes section).

Genome copies/day/person
Green solid line
Log₁₀ scale

Genome copies/L x Daily wastewater influent (L) = genome copies/day
 Genome copies/day ÷ population = genome copies/day/person




Rolling seven-day average of **new cases** in the catchment
 Grey dotted line (shaded)
 Linear Scale
 This data is not available for all sites, and subject to change.


Note the scales may differ for each graph.
 Wastewater results are on log₁₀ scale, while case numbers are on a linear scale.

Northland


Whangarei




Paihia




Kaiwaka





Mangawhai



Kaikohe




Dargaville




Status ● Detected ● Not detected


Kaitiaki




Kawakawa



Hikurangi




Ruakaka




Auckland


Auckland East




Auckland Southwest




Army Bay





Wellsford




Auckland (Combined)




Auckland West




Omaha




Snells/Algies




Beachlands




North Shore




Pukekohe




Clarks Beach




Warkworth



Waiuku




Kawakawa Bay




Waikato


Whangamata




Thames




Te Awamutu





Whitianga




Hamilton



Cambridge




Taupo




Bay of Plenty and Gisborne


Maketu




Gisborne




Mt Maunganui





Te Puke








Tauranga



Rotorua




Status ● Detected ● Not detected




Hawke's Bay


Hastings




Waipawa




Otane





Napier



Takapau




Waipukurau




Status ● Detected ● Not detected

Wairoa




Taranaki


Eltham




Opunake




New Plymouth





Patea



Hawera




Stratford




Status ● Detected ● Not detected


Kaponga



Manaia




Waverley




Manawatu-Wanganui


Levin




Palmerston North




Whanganui Urban





Feilding




Taumarunui




Pahiatua




Dannevirke



Woodville




Eketahuna




Wellington


Hutt Valley




Wellington (Western)




Porirua



Paraparaumu




Wellington (Moa Point)




Status ● Detected ● Not detected


Martinborough




Carterton




Masterton




Otaki



Featherston




Greytown




Tasman, Nelson, and Marlborough


Picton




Motueka




Nelson Central/North



Blenheim




Richmond/Nelson South




Status ● Detected ● Not detected

West Coast and Canterbury


Timaru




Rolleston & Eastern Selwyn




Leeston





Christchurch



Ashburton




Oamaru




Status ● Detected ● Not detected


Reefton




Westport




Hokitika




Rangiora




Kaikoura




Greymouth




Kaiapoi



Amberley




Woodend




Otago and Southland


Dunedin (Mosgiel)




Dunedin (Green Island)




Invercargill





Bluff



Queenstown




Dunedin (Tahuna)




Status ● Detected ● Not detected


Alexandra




Cromwell




Balclutha



Gore



Wanaka



Region/Site	2021																					2022																			
	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13	14						
Dunedin (Tahuna)																																									
Milton																																									
Mosgiel																																									
Oamaru																																									
Queenstown																																									
Wanaka																																									
Southland																																									
Bluff																																									
Gore																																									
Invercargill																																									

*PS and MH refer to Pump Station and Manhole sites

Acknowledgements

This work represents the combined efforts of a large number of individuals and organisations.

We continue to be indebted to the teams across the country who are collecting the wastewater that underpins this work.

The wastewater analysis has been undertaken at the ESR Kenepuru (in Porirua) and Christchurch laboratories by a team which may on any given week include contributions from: Kirsten Carter, Joanne Chapman, Dawn Croucher, Joanne Hewitt, Joycelyn Ho, Anower Javed, Susan Lin, Olivia Macrae, Ashley McDonald, Andrew Ng, Ashley Orton, Paula Scholes and Fatiha Sulthana. Data science analysis, visualization and reporting is the result of team effort from: Franco Andrews, Bridget Armstrong, Raewyn Campbell, Gerhard de Beer, Richard Dean, Brent Gilpin, Joanne Hewitt, Dawen Li, Helen Morris and Bindu Priya. Ongoing support for this work from the Ministry of Health and ESR management is appreciated.

Notes

Sites and frequency of sample collection: The catchment population sites selected for the surveillance range from approximately 100 to over 1,000,000 individuals. The sites cover all regions of the country. Most major towns and all cities, as well as many smaller communities, are included. In early 2022, the wastewater catchment areas cover over 80% of the population connected to wastewater treatment plants. The sites from which samples have been collected have varied over the last 12 months. New sites may be added over time, and/or sampling may reduce in frequency or cease for other sites. The selection and frequency of sampling vary depending on the local population, access to wastewater collection points, staff availability to collect samples and risk factors. When included, samples are collected at least weekly, with twice weekly sampling being common. A number of samples have also been collected from non-WWTP sites (manholes and pump stations- mostly in Auckland).

Sampling method: The preferred option is to automatically collect a 24 hour 'composite' sample. This is where a pump automatically collects a small volume of wastewater every 15 minutes over 24 hours using a composite sampler. These samplers are available in some wastewater treatment plants. When composite samplers are not available, 'grab' samples are collected. These range from a sample being taken at a single point in time, to 3 samples taken over 30 minutes, to samples collected over a day. Grab samples represent only the composition of the source at that time of collection and may not be as representative as a 24 hour composite sampler. More variation may be expected with grab samples.

Laboratory analysis of wastewater samples: Samples are sent from each wastewater treatment plant to one of the ESR laboratories (Porirua or Christchurch). Processing of each sample commences within an hour or two of receipt. Processing involves the concentration of virus from 250 mL sample to approx. 1 mL using centrifugation and polyethylene glycol. Viral RNA is then extracted from a small volume of 0.2 mL concentrate to give a final volume of 0.05 mL. The presence of SARS-CoV-2 RNA is determined using RT-qPCR. SARS-CoV-2 is considered detected when any of the RT-qPCR replicates are positive.

RT-qPCR: Reverse transcription (RT) to convert RNA to complementary DNA (cDNA), followed by quantitative PCR (qPCR). RT-qPCR is used for detection and quantification of viral RNA.

Method sensitivity: The protocol used to concentrate SARS-CoV-2 from wastewater allows for the sensitive detection of SARS-CoV-2 by RT-qPCR. ESR has shown that when 10 individuals are actively shedding SARS-CoV-2 RNA in a catchment of 100,000 individuals, there was a high likelihood of detecting viral RNA in wastewater (<https://doi.org/10.1016/j.watres.2021.118032>). Shedding by one individual may be detected in wastewater, but it does depend on many factors including the amount and duration of shedding. Very low levels in wastewater may be not able to be quantified (i.e., less than the limit of quantification- see below).

SARS-CoV-2 RNA detected (positive result): A positive detection in the wastewater indicates that at least one person has been shedding SARS-CoV-2 into the wastewater at some point during the time period that the sample was being collected. In some cases, detections could also be due to the shedding of low levels of SARS-CoV-2 RNA by a recently recovered case. The detection of SARS-CoV-2 RNA does not indicate that infectious virus is present.

SARS-CoV-2 RNA not detected (negative result): A negative result can occur because there are no active 'shedding' cases in the catchment or because the SARS-CoV-2 RNA concentration is too low to be detected, most likely because there are a very low number of cases in the wastewater catchment. Therefore, negative finding does not necessarily guarantee the absence of COVID-19 in the community.

Viral loads and normalisation: When detected, the SARS-CoV-2 RNA concentration is calculated as genome copies per mL or per L wastewater. This is then converted to a viral load of **genome copies/day/person**. This conversion takes into account the flow rate of wastewater entering the treatment plant (the influent) and the population in the catchment. The **flow rate** is the total volume (m³ per day) recorded at the inlet of the wastewater treatment plant over 24 hours. This is a **population-normalised viral load**. Currently, the flow rate is the average annual flow rate, but will be replaced with daily flow rate when available (note that rainfall may significantly increase the flow rate at the inlet, diluting the sample, and may result in lower concentrations and a false negative result).

In future, SARS-CoV-2 RNA concentrations will also be normalised by testing for the presence of pepper mild mottled virus (PMMoV). PMMoV is a virus that infects peppers but not humans. Consumption of peppers or pepper products, such as chilli sauce, means that PMMoV is detected in wastewater – normally at very high concentrations. Therefore, PMMoV has been found to be a useful proxy for the amount of faecal material in a wastewater sample. For normalisation, the concentration of SARS-CoV-2 RNA is divided by that of PMMoV in each sample. Different normalisation methods may result in changes to some data points but trends are unlikely to change significantly.

Limit of quantification: The lowest concentration of the target that can be reliably quantified is referred to as the limit of quantification. For those samples where SARS-CoV-2 is detected but cannot be quantified, a value of 5 genome copies/mL wastewater is used. While a standard method is being used, virus recovery can vary from sample to sample, and this may affect the quantitation.

Data subject to change: Data generated for the New Zealand Wastewater COVID-19 Surveillance Programme should be considered provisional and may be subject to change. Data may be incomplete for the most recent 2-week period due to processing, testing and reporting delays.

Data not shown:

- Data from 'ad hoc' sampling locations including from individual facilities/building (e.g., workplaces, prisons, MIQs) are not included.
- Results from certain samples may not be shown, as the result was either deemed invalid, or the sample could not be tested (e.g., leaked in transit, not labelled).